

Mycolic Acid-Containing Actinomycetes Associated with Activated Sludge Foam

Chi Nam Seong*, Young Sook Kim, Keun Shik Baik, Soon Dong Lee¹,
Yung Chil Hah¹, Seung Bum Kim², and Michael Goodfellow²

Department of Biology, College of Natural Sciences,
Suncheon National University, Suncheon 540-742;

¹Department of Microbiology, College of Natural Sciences, and Research Center for
Molecular Microbiology, Seoul National University, Seoul 151-742, Korea;

²Department of Agricultural and Environmental Science, University of Newcastle,
Newcastle upon Tyne, NE1 7RU, U.K.

(Received April 22, 1999 / Accepted May 31, 1999)

Mycolic acid-containing actinomycetes associated with extensive foaming in the aeration basin of the activated sludge process were isolated and analyzed by phenotypical, chemotaxonomical and phylogenetic methods. Whole cell sugar patterns of two isolates were pattern A. The nearly complete sequences of the 16S rRNA genes (rDNAs) of the isolates were determined and compared by using several tree-making algorithms. With polyphasic methods, strain SCNU1 was identified as *Gordona sputi*, and strain SCNU5 assigned to the genus *Tsukamurella*. The presence of opportunistic pathogens of chronic lung infections within foams can cause public health problems and render waste-treatment processes inefficient.

Key words: Mycolic acid, foaming, *Gordona sputi*, *Tsukamurella*

Bulking and foaming are the most common problems in the activated sludge process. Bulking is a macrostructure failure due to the presence of large numbers of filamentous microorganisms in activated sludge. These organisms interfere with the compaction and settling of activated sludge either by producing a very diffuse floc structure or by growing in profusion beyond the confines of the activated sludge floc into the bulk medium thereby causing bridging between flocs (1, 26).

The formation of scum or foam in the activated sludge process is also caused by filamentous bacteria, notably *Microthrix parvicella*, and mycolic acid containing actinomycetes. *Nocardia* sp., *Gordona* sp., *Tsukamurella* sp., *Actinomadura* sp., *Micromonospora* sp., *Mycobacterium* sp., and some *Streptomyces* sp. are known to produce foam in activated sludge (1, 2, 6, 25). These bacteria have hydrophobic cell surfaces and when present in sufficient numbers in activated sludge, they render flocs hydrophobic and hence amenable to the attachment of air-bubbles. The air bubble-floc aggregate is less dense than water and hence floats to the surface of the sludge. The hydrophobic flocs tend to stay at the surface of

the sludge where they accumulate to form a thick, chocolate-brown coloured foam or scum. Activated sludge foams cause a number of problems, namely extra-house keeping on the part of the operator; blockage of scum removal systems; reduction of oxygen transfer at the surface of mechanically aerated basins; carriage and dispersal of pathogens in wind blown scum; drying of scum with resultant cleaning and possibly odour problems, and reduction of effluent quality through an increase in effluent suspended solids and biological oxygen demand if the scum reaches the final effluent (26).

The primary aim of the present study was to isolate, characterize and identify mycolic acid-containing actinomycetes associated with extensive foaming in the aeration basin. First, the chemical composition and physiological properties of the strains were investigated. Then, the almost complete sequences of the 16S rRNA genes (rDNAs) of isolates were determined and compared by using several tree-making algorithms (11, 13, 18).

Materials and Methods

Isolation of actinomycetes associated with foam

The foams used in this study were obtained from

* To whom correspondence should be addressed.
(Tel) 82-661-750-3613; (Fax) 82-661-750-3608
(E-mail) scnu@suncheon.suncheon.ac.kr

three full-scale plants in Sunchon, Korea. Two procedures such as micromanipulation and direct plating were applied to isolate the foam-forming strains. Approximately 0.5 mg of foams were diluted with 10 ml of distilled water. The mixture was stirred in the tube mixture for 20 minutes and the flocs were allowed to settle. Then, an aliquot (0.5 ml) of the supernatant obtained from the surface was used as an inoculum. First, supernatant was smeared on the sloppy agar (0.5%, w/v) slice, and the microorganisms on the agar slice was picked with a micromanipulator (Narishigae) attached to an inverted microscope (Nikon). Otherwise, a direct plating technique was used to isolate actinomycetes. Several agar media were used to isolate the actinomycetes. R2A (22), Tween media (BSM, PRM-1, EPM-1, EPM-2, and NTM; 16), SCY, I, Sludge hydrolysate (6), Tryptone-yeast extract-glucose agar (TYGA;10) and MP (25) medium were prepared as described previously. Inoculated plates were incubated for 3 weeks at 28°C. The colonies were examined daily using a microscope. For the subsequent cultural work, glucose yeast extract agar (GYEA) medium and the same incubation conditions were used. Isolated strains were maintained as mycelial fragment suspensions (20% [v/v] glycerol) at -30°C.

Morphology

Mycelial fragmentation, aerial mycelium color, pigmentation and colony morphology were observed on GYEA, oatmeal agar (ISP3), and inorganic salts-starch casein agar (ISP4) medium after incubation at 30°C daily for 7 days (24). Melanin production was observed on peptone-yeast extract-iron agar (ISP6) and tyrosine agar (ISP7).

Mycolic acid determination

Biomasses were obtained from liquid cultures in Sauton's broth medium (Difco) at 28°C for 5 days, freeze dried, and kept refrigerated for further analysis. Extraction of mycolic acids was performed using alkaline methanolysis (20). Thin layer chromatography, using Kieselgel 60 Aluminium plate (Merck), was used to detect the mycolic acids. A single development with petroleum ether (bp 60~80°C)-toluene (70:30, v/v) was used. Detection was performed with a 5% ethanolic molybdophosphoric acid spray followed by heating at 180°C for 5 min.

Menaquinone analysis

About 50 mg of dried biomasses was treated with chloroform/methanol (2:1, v/v) by shaking overnight. The extracts were concentrated in vacuo using an electric aspirator and the cell debris were removed by centrifugation. The solvents were evaporated

completely, and the remnants were resuspended in hexane. High performance liquid chromatographic separation of the quinones was done with an ODS Hypersil column (2004.6 mm, particle size 5 µm, Hewlett Packard) and acetonitrile/tetrahydrofuran (70:30, v/v) as the mobile phase. The flow rate was 1 ml/min at 37°C, and detector was a UV detector operated at 254 nm.

Whole cell sugar analysis

Whole cell sugars extracted as alditol acetates (7) were analyzed using a gas chromatograph (Hewlett Packard 5890A) fitted with a flame ionization detector. Separation was achieved using a 0.53 mm × 30 m SP 2380 (Supelco) fused silica capillary column. The temperature was programmed to hold at 160°C for 2 min, then to rise by 5°C/min. The Injector temperature was held at 250°C, and the detector at 300°C.

Phenotypic tests

The strains were examined for 83 unit characters including pH- and temperature-dependent growth, enzyme synthesis, resistance to antibiotics and chemicals, antibiosis, utilization of carbon and nitrogen sources, and acid production from sugars (10, 29).

DNA extraction and amplification of 16S rDNA

Chromosomal DNA was isolated from the test strains using a procedure (3) slightly modified from that of Pitcher *et al.* (21). PCR amplification of 16S rDNA was performed using a capillary Rapidcyclor (Idaho Technology). The following conditions were used for amplification using primers 27f and 1525r: denaturation at 94°C for 1 min, annealing at 55°C for 1 min and elongation at 72°C for 3 min for each cycle. A total of 30 cycles were performed followed by a final elongation step at 72°C for 10 min and cooling at 25°C for 1 min.

Sequencing of 16S rDNA

The amplified fragments were directly sequenced using a Taq DyeDeoxy terminator Cycle Sequencing Kit (Applied Biosystems) and previously described oligonucleotide primers (3). Sequencing gel electrophoresis was carried out and nucleotide sequences were automatically obtained using an Applied Biosystems DNA sequencer (model 373A) with software provided by the manufacturer.

Phylogenetic analysis

The two 16S rDNA sequences were aligned manually with mycolic acid containing actinomycete

Table 1. Chemotaxonomic characteristics of strain SCNU1 and SCNU5

Strain	Wall diamino acid	Wall sugars	Wall chemotype ^a	Major menaquinones
SCNU1	<i>meso</i> -DAP	Galactose, arabinose	IV	MK-8(H ₂), MK-9(H ₂)
SCNU5	<i>meso</i> -DAP	Galactose, arabinose	IV	MK-9(H ₀)

^a Wall chemotype as described by Lechevalier and Lechevalier(17).

nucleotide sequences derived from the Ribosomal Database Project (19) and EMBL/GenBank database using the AL 16S program (4). The reference sequences had the following accession numbers: Z46753 (*Corynebacterium blutamicum*), X79290 (*Dietzia maris*), X80633 (*Gordona aichiensis*), X80635 (*G. amarae*), X79287 (*G. bronchialis*), X80632 (*G. rubropertincta*), X80634 (*G. sputi*), X79286 (*G. terrae*), X52917 (*Mycobacterium tuberculosis*), X57949 (*Nocardia asteroides* DSM43005), X79288 (*Rhodococcus rhodochrous*), Z46751 (*Tsukamurella paurometabola*), X87340 (*G. hydrophobica*), X93485 (*G. hirsuta*), Z35435 (*Skermania piniformis*), X92981 (*T. pulmonis*), and Y12246 (*T. tyrosinosolvens*). 16S rDNA sequences of strain SCNU1 and SCNU5 were deposited in the GenBank under accession number AF150493 and AF150494, respectively. Evolutionary trees for the datasets were inferred by using three algorithms, namely the neighbour-joining (23), least squares (19) and maximum-parsimony (15) methods. Evolutionary distance matrices for the neighbour-joining and least-squares methods were generated as described by Jukes and Cantor (12). The PHYLIP package (8) was used for making trees with the three algorithms. The resultant unrooted tree topologies were evaluated by carrying out bootstrap analysis of the neighbour-joining method data based on 1000 re-samplings using the SEQBOOT and CONSENSE programs in the PHYLIP package.

Results

Estimation of isolation medium

Several agar media were used for the isolation of actinomycetes associated with activated sludge foam. Most colonies which developed on the media were non-actinomycete bacteria. While large numbers of filamentous and mycelial fragments were observed microscopically in the foam and diluent, the recovery rate was relatively poor on the agar plate. Actinomycetes were recovered only on I and TYGA medium, while other media were not suitable for the growth of actinomycetes.

Chemotaxonomy

Two mycolic acid-containing strains were pure

cultured. They contained *meso*-diaminopimelic acid (*meso*-DAP). TLC analysis of mycolic acids confirmed that strain SCNU1 and SCNU5 belong to the genus *Gordona*, and *Tsukamurella*, respectively. Characteristic sugars and major menaquinone composition of strains were shown in Table 1. The two strains contained the same whole cell sugars such as arabinose and galactose (whole cell sugar pattern A). Strain SCNU5 contained menaquinone of 9 isoprene units, designated as MK-9, without hydrogenation (MK-9(H₀)), while strain SCNU1 had menaquinones of different isoprene units but the same degree of hydrogenation, designated as MK-8 (H₂) and MK-9 (H₂).

Phenotypic properties

The properties of the strains were described in Table 2. The strains did not produce aerial mycelium and diffusible pigment. They could grow both at pH 4 and 9. They did not produce melanin. They all degraded gelatin and esculin, but not starch, casein, elastin and xanthine. They were sensitive to streptomycin, gentamicin, and vancomycin, but resistant to bacitracin. The strains utilized succinate, pyruvate, and propionate, but not lactate and citrate. They did not produce acid from most of the sugars tested. The strains did not show antimicrobial activity.

Phylogenetic analysis

The almost complete 16S rDNA sequences of the strains were compared with the corresponding nucleotide sequences of mycolic acid containing actinomycetes by using three tree-making algorithms (i.e. the neighbour-joining, least squares and maximum-parsimony as shown in Fig. 1). 16S rDNA of strain SCNU1 was identical to that of *G. sputi* except for one nucleotide. Nucleotide sequences of strain SCNU5 showed 99.73% similarity with *T. pulmonis* and 99.66% with *T. tyrosinosolvens* (Table 3).

Comparison of *Tsukamurella* sp. strain SCNU5 with related species

The isolate SCNU5 was compared with the most related species, *T. pulmonis* and *T. tyrosinosolvens*. Physiologically, SCNU5 differs from *T. pulmonis* in i) its inability to degrade urea, ii) its ability to liquify gelatin, iii) its sensitivity to

Table 2. Phenotypic properties of the mycolic acid-containing actinomycete strain SCNU1 and SCNU5 isolated from activated sludge foam

Unit characters	SCNU1	SCNU5	Unit characters	SCNU1	SCNU5
Substrate mycelium color	Pink	—	D-Melibiose	—	—
Melanin production	—	—	Cellobiose	—	—
Degradation of:			Maltose	—	—
Hypoxanthine	—	—	Trehalose	—	—
Elastin	—	—	Mannitol	—	+
Xanthine	—	—	Salicin	—	—
Casein	—	—	Arabinose	—	—
Urea	+	—	Adonitol	—	—
Starch	—	—	Melezitose	—	—
Tyrosine	—	—	Inositol	—	—
Guanine	—	—	Xylose	—	—
Gelatin	+	+	Fructose	—	—
Esculin	+	+	Mannose	—	—
Growth in the presence of (%):			Sucrose	—	—
NaCl (4)	+	+	Erythritol	—	+
NaCl (7)	+	—	Galactose	—	—
NaCl (10)	+	—	D-Sorbitol	—	+
Sodium azide (0.01)	+	—	Glucose	—	—
Sodium azide (0.02)	+	—	Dulcitol	—	—
Phenol (0.1)	—	—	Xylitol	—	—
Phenylethanol (0.1)	+	—	Dextrin	—	—
Phenylethanol (0.3)	—	—	Utilization of nitrogen sources:		
Thallos acetate (0.001)	+	—	DL- α -Amino-n-butyric acid	—	—
Thallos acetate (0.01)	+	—	L-Cysteine	—	+
Crystal violet (0.0001)	+	+	L-Valine	—	+
Resistance to antibiotics (μ g/ml):			L-Hydroxyproline	—	—
Rifampicin (50)	+	—	L-Histidine	—	—
Penicillin G (10 i.u.)	—	+	L-Phenylalanine	—	+
Streptomycin (10)	—	—	Antimicrobial activity against:		
Gentamicin (10)	—	—	<i>A. niger</i>	—	—
Bacitracin (10 i.u.)	+	+	<i>B. subtilis</i>	—	—
Vancomycin (30)	—	—	<i>C. albicans</i>	—	—
Neomycin (30)	—	—	<i>E. coli</i>	—	—
Utilization of organic acid:			<i>P. fluorescence</i>	—	—
Oxalate	—	—	<i>S. cerevisiae</i>	—	—
Succinic acid	+	+	Enzyme synthesis:		
Pyruvic acid	+	+	Lipolysis	—	—
Lactic acid	—	—	Pectin hydrolysis	—	—
Mucic acid	—	—	Nitrate reduction	+	—
Propionic acid	+	+	Production of hydrogen sulfide	+	+
Gluconic acid	—	—	Growth at:		
Citric acid	—	—	pH 4	+	+
Acid production from sugars:			pH 9	+	+
Lactose	—	—	10°C	+	—
L-Rhamnose	—	—	37°C	+	+
Raffinose	—	—	45°C	—	—

rifampicin and streptomycin, and iv) its ability to utilize gluconic acid. SCNU5 differs from *T. tyrosinosolvens* in i) its inability to degrade hypoxanthine and xanthine, ii) its ability to liquify gelatin, and iii) its ability to utilize citric acid.

Discussion

For the success of selective isolation of foam-asso-

ciated microorganisms, isolation procedures and selective media were tested. Only two media, I and TYGA were found to be adequate for the isolation of actinomycetes. Using the micromanipulation procedure, few filamentous microorganisms were recovered. Finally, two mycolic acid-containing actinomycetes were isolated. Further studies on the choice of isolation medium corresponding to the status of sludge composition and operation style should be carried out.




Fig. 1. A neighbour-joining tree (23) based on nearly complete 16S rRNA sequences of mycolic acid-containing bacteria and representative species of related taxa. The scale bar indicates 0.01 nucleotide substitutions per nucleotide position.

In this study, we isolated two mycolic acid-containing actinomycetes, then analyzed and identified them with different categories of phenotypic, chemotaxonomic and phylogenetic methods. Two isolates were different each other in their mycolic acid profile. Whole cell sugar and menaquinone composition were determined by the chromatographic method. They all contained arabinose and galactose as characteristic whole cell sugars. Thus, the two strains were classified as wall chemotype IV. In contrast, their major menaquinones were different each other. From the chemotaxonomical

Table 4. Differential physiological characteristics of strain SCNU5, *T. pulmonis* and *T. tyrosinosolvens*

Unit characters	SCNU5	<i>T. pulmonis</i> ^a	<i>T. tyrosinosolvens</i> ^b
Diffusible pigment	–	–	–
Degradation of:			
Hypoxanthine	–	–	+
Elastin	–	–	N
Xanthine	–	–	+
Casein	–	–	N
Urea	–	+	N
Tyrosine	–	–	+
Guanine	–	–	N
Gelatin	+	–	–
Esculin	+	+	N
Resistance to antibiotics (µg/l):			
Rifampicin (50)	–	+	N
Streptomycin (10)	–	+	N
Utilization of organic acid:			
Lactic acid	–	–	–
Gluconic acid	–	+	N
Citric acid	–	–	+
Enzymatic activity:			
Nitrate reduction	–	–	N
Growth at:			
37°C	+	+	N
45°C	–	–	–

Source of data from ^aYassin *et al.* (30) and ^bYassin *et al.* (31). N, data not available.

analysis, strain SCNU1 and SCNU5 were assigned to the genus *Gordona* and *Tsukamurella*, respectively.

Properties of the isolates were determined against 83 unit characters including morphology, degradation, pH- and temperature-dependent growth, enzyme synthesis, resistance to antibiotics and chemicals, antibiosis, utilization of carbon and nitrogen sources, and acid production from sugars. The two strains did not produce acid from most organic acids. In addition, their antimicrobial activity was not found.

The almost complete sequences of the 16S rRNA genes (rDNA) of two strains were determined and compared by several tree-making algorithms. Strain SCNU1 was almost identical to *G. sputi*. 16S rDNA

Table 3. Levels of 16S rDNA similarity between the strain SCNU5 and related species

Microorganism	% 16S rDNA similarity					
	SCNU5	<i>T. spumae</i>	<i>T. wratislaviensis</i>	<i>T. inchoensis</i>	<i>T. pulmonis</i>	<i>T. paurometabola</i>
<i>T. spumae</i>	99.66					
<i>T. wratislaviensis</i>	95.79	95.72				
<i>T. inchoensis</i>	99.12	99.52	95.72			
<i>T. pulmonis</i>	99.73	99.52	95.88	99.25		
<i>T. paurometabola</i>	99.18	99.39	95.85	99.46	99.25	
<i>T. tyrosinosolvens</i>	99.66	99.45	95.95	99.38	99.73	99.38

nucleotide sequence of SCNU5 differed with that of *T. pulmonis* and *T. tyrosinosolvens* by 4 and 5 nucleotides, respectively.

The similarity levels between species in the genus *Tsukamurella* were known to be relatively high (30, 31). In contrast to 16S rRNA analysis, the physiological tests revealed some differences between strain SCNU5 and the other two strains. Thus, the SCNU5 strain can be placed in a new species of the genus *Tsukamurella*. For the complete identification of strain SCNU5, molecular fingerprinting techniques, such as random amplified polymorphic DNA (RAPD) PCR and DNA-DNA hybridization, should be carried out.

G. amarae have been frequently isolated from activated sludge foam in previous studies (27). In contrast, *G. sputi* associated with sludge foam was not yet reported. In particular, *Gordona sputi* and a number of species in the genus *Tsukamurella* are infectious agents, and have been commonly isolated from the sputum of patients with chronic lung infections or from the blood of patients with cardiac pacemaker implants (27, 30, 31). The isolation of *G. sputi* in this study confirmed the existence of bacterial pathogens in activated sludge foam. Based on these results mycolic acid-containing actinomycetes can not only make the waste-treatment process inefficient by foaming, but also render the pathogens to wind dispersal.

Acknowledgment

The authors wish to acknowledge the financial support of the Korea Research Foundation made in the program year of 1997.

References

- Blackall, L.L., J.H. Parlett, A.C. Hayward, D.E. Minnikin, P.F. Greenfield, and A.E. Harbers. 1989. *Nocardia pinensis* sp. nov., an actinomycete found in activated sludge foams in Australia. *J. Gen. Microbiol.* **135**, 1547-1558.
- Blackhall, L.L., E.M. Seviour, M. Cunningham, R.J. Seviour, and P. Hugenholtz. 1994. "*Microthrix parvicella*" is a novel, deep branching member of the actinomycetes subphylum. *Syst. Appl. Microbiol.* **17**, 513-518.
- Chun, J. and M. Goodfellow. 1995. A phylogenetic analysis of the genus *Nocardia* With 16S rRNA gene sequences. *Int. J. Syst. Bacteriol.* **45**, 240-245.
- Chun, J. 1995. Computer-assisted classification and identification of actinomycetes. PhD thesis, University of Newcastle upon Tyne.
- Chun, J., L.L. Blackall, S.O. Kang, Y.C. Hah, and M. Goodfellow. 1997. A proposal to reclassify *Nocardia pinensis* Blackall et al. as *Skermania pniformis* gen. nov., comb. nov. *Int. J. Syst. Bacteriol.* **47**, 127-131.
- Eikelboom, D.H. 1994. The *Microthrix parvicella* puzzel. *Water Science Technol.* **29**, 271-279.
- Englyst, H.N., and J.H. Cummings. 1984. Simplified method for the measurement of total non-starch polysaccharides by gas-chromatography of constituent sugars as alditol acetates. *Analyst.* **101**, 937-942.
- Felsenstein, J. 1993. PHYLIP (Phylogeny Inference Package) version 3.5.1. Department of Genetics, University of Washington, Seattle, USA.
- Fitch, W.M. and E. Margoliash. 1967. Construction of phylogenetic trees: a method based on mutation distances as estimated from cytochrome c sequences is of general applicability. *Science* **155**, 279-284.
- Gordon R.E., D. A. Barnett, J.E. Handerhan, and C.H. Pang. 1974. *Nocardia coeliaca*, *Nocardia autotrophica*, and the *Nocardia* Strain. *Int. J. Syst. bacteriol.* **24**, 54-63.
- Huelsenbeck, J.P. and D.M. Hillis. 1993. Success of phylogenetic method in the four-taxon case. *Syst. Biol.* **42**, 247-264.
- Jukes, T.H. and C.R. Cantor. 1969. Evolution of protein molecules. In *Mammalian Protein Metabolism*, vol. 3, pp. 21-132. Edited by H.N. Munro. New York; Academic Press.
- Kim, J. 1993. Improving the accuracy of phylogenetic estimation by combining different methods. *Syst. Biol.* **42**, 331-340.
- Kim, S.B., M.Y. Kim, C.N. Seong, and Y.C. Hah. 1996. Lipid analysis of streptomycetes isolated from volcanic soil. *J. Microbiol.* **34**, 184-191.
- Kluge, A.G. and F.S. Farris. 1969. Quantitative phyletics and the evolution of anurans. *Syst. Zool.* **18**, 1-32.
- Kocianova, E., C.F. Forster, and R.J. Foot. 1994. What is *Microthrix parvicella*? *J. Appl. Bacteriol.* **76**, 301-306.
- Lechevalier, M.P. and M.P. Lechevalier. 1970. Chemical composition as a criterion in the classification of aerobic actinomycetes. *Int. J. Syst. Bacteriol.* **20**, 435-443.
- Li, W.H. and M. Gony. 1990. Statistical tests of molecular phylogenies. *Methods Enzymol.* **183**, 645-659.
- Maidak, B.L., G.J. Olsen, N. Larsen, R. Overbeek, M.J. McCaughey, and C.R. Woese. 1996. The Ribosomal Database Project (RDP). *Nucleic acids Res.* **24**, 82-85.
- Minnikin, D.E., I.G. Hutchinson, A.B. Caldicott, and M. Goodfellow. 1980. Thin-layer chromatography of methanolysates of mycolic acid-containing bacteria. *J. Chromatogr.* **188**, 221-233.
- Pitcher, D.G., N.A. Saunders, and R.J. Owen. 1989. Rapid extraction of bacterial genomic DNA with guanidium thiocyanate. *Lett. Appl. Microbiol.* **8**, 151-156.
- Reasoner, D.J. and E.E. Geldreich. 1985. A new medium for the enumeration and subculture of bacteria from potable water. *Appl. Environ. Microbiol.* **49**, 1-7.
- Saitou, N. and M. Nei. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**, 406-425.
- Shirling, E.B. and D. Gottlieb. 1966. Methods for characterization of *Streptomyces* species. *Int. J. Syst. Bacteriol.* **16**, 313-340.
- Slijkhuis, H. 1983. *Microthrix parvicella*, a filamentous bacterium isolated from activated sludge: cultivation in a chemically defined medium. *Appl. Environ. Microbiol.* **46**, 832-839.
- Soddell, J.A. and R.J. Seviour. 1990. Microbiology of

- foaming in activated sludge plants. *J. Appl. Bacteriol.* **65**, 145-176.
27. **Steingrube, V.A. R.W. Wilson, B.A. Brown, K.C. Brown, Z. Blacklock, J.L. Gibson, and R.J. Wallace.** 1997. Rapid identification of clinically significant species and taxa of aerobic actinomycetes, including *Actinomadura*, *Gordona*, *Nocardia*, *Rhodococcus*, *Streptomyces*, and *Tsukamurella* isolates, by DNA amplification and restriction endonuclease analysis. *J. Clin. Microbiol.* **35**, 817-822.
28. **Van Veen, W.L.** 1973. Bacteriology of activated sludge, in particular the filamentous bacteria. *Antonie van Leeuwenhoek* **39**, 189-205.
29. **Williams, S. T., M. Goodfellow, G. Alderson, E.M.H. Wellington, P.H.A. Sneath, M.J. Sackin.** 1983. Numerical classification of *Streptomyces* and related genera. *J. Gen. Microbiol.* **129**, 1743-1813.
30. **Yassin, A.F., F.A. Rainey, H. Brzezinka, J. Burghardt, P. Seifert, K. Feldmann, and K.P. Schaal.** 1996. *Tsukamurella pulmonis* sp. nov. *Int. J. Syst. Bacteriol.* **46**, 429-436.
31. **Yassin, A.F., F.A. Rainey, J. Burghardt, H. Brzezinka, S. Schmitt, P. Seifert, O. Zimmermann, H. Mauch, D. Gierth, I. Lux, and K.P. Schaal.** 1997. *Tsukamurella tyrosinosolvens* sp. nov. *Int. J. Syst. Bacteriol.* **47**, 607-614.