

Enhancing the Intrinsic Bioremediation of PAH-Contaminated Anoxic Estuarine Sediments with Biostimulating Agents

Quang-Dung Bach, Sang-Jin Kim¹, Sung-Chan Choi² and Young-Sook Oh*

Department of Environmental Engineering and Biotechnology, Myongji University, Yongin 449-728, Republic of Korea

¹Microbiology Lab., Korea Ocean Research and Development Institute, P.O. Box 29, Ansan 425-600, Republic of Korea

²Department of Environmental Sciences and Biotechnology, Hallym University, Chunchon 200-702, Republic of Korea

(Received April 18, 2005 / Accepted June 7, 2005)

Estuarine sediments are frequently polluted with hydrocarbons from fuel spills and industrial wastes. Polycyclic aromatic hydrocarbons (PAHs) are components of these contaminants that tend to accumulate in the sediment due to their low aqueous solubility, low volatility, and high affinity for particulate matter. The toxic, recalcitrant, mutagenic, and carcinogenic nature of these compounds may require aggressive treatment to remediate polluted sites effectively. In petroleum-contaminated sediments near a petrochemical industry in Gwangyang Bay, Korea, *in situ* PAH concentrations ranged from 10 to 2,900 µg/kg dry sediment. To enhance the biodegradation rate of PAHs under anaerobic conditions, sediment samples were amended with biostimulating agents alone or in combination: nitrogen and phosphorus in the form of slow-release fertilizer (SRF), lactate, yeast extract (YE), and Tween 80. When added to the sediment individually, all tested agents enhanced the degradation of PAHs, including naphthalene, acenaphthene, anthracene, fluorene, phenanthrene, fluoranthene, pyrene, chrysene, and benzo[a]pyrene. Moreover, the combination of SRF, Tween 80, and lactate increased the PAH degradation rate 1.2-8.2 times above that of untreated sediment (0.01-10 µg PAH/kg dry sediment/day). Our results indicated that *in situ* contaminant PAHs in anoxic sediment, including high molecular weight PAHs, were degraded biologically and that the addition of stimulators increased the biodegradation potential of the intrinsic microbial populations. Our results will contribute to the development of new strategies for *in situ* treatment of PAH-contaminated anoxic sediments.

Key words: biodegradation, bioremediation, anoxic sediment, PAH, biostimulation

Polycyclic aromatic hydrocarbons (PAHs) are artificial fused-ring compounds that enter sediment systems from coke and petroleum-refining industries, accidental spills and leakages, and rainwater runoff from roadways (Kanaly and Harayahama, 2000; Karthikeyan and Bhandari, 2001; Wang *et al.*, 2001; Rothermich *et al.*, 2002; Chang *et al.*, 2003). PAHs pose an environmental threat primarily because of their toxicity, low volatility, resistance to microbial degradation, and ability to sorb to sediments. Documented impacts on critical habitats include contamination of benthic ecosystems, with subsequent harm to the marine food chain (Coates *et al.*, 1997; Elliott, 2001; Wang *et al.*, 2001).

Researchers have studied the anaerobic biodegradation of PAHs in sediment since the late 1980s (Lovley *et al.*, 1989; Coates *et al.*, 1997; Hayes *et al.*, 1999; Rothermich *et al.*, 2002). For anaerobic processes, PAH degradation coupled to sulfate reduction is the most relevant to coastal

marine sediments, because sulfate is abundant in seawater, whereas nitrate concentrations are typically low and Fe(III) is often only slightly available (Coates *et al.*, 1997; Rothermich *et al.*, 2002). Apart from a study on the methanogenic degradation of benzene (Weiner and Lovley, 1998), few reports on the methanogenic transformation of PAHs have been published to our knowledge.

PAHs are generally slowly degraded in the environment, particularly in marine systems (Kanaly and Harayahama, 2000; Nieman *et al.*, 2001; Han *et al.*, 2003). The low degradation rates of PAHs may be due to their stable structures, adsorption onto particulate matter, or nutrient deficits in the environment (Gibbons, 1991; Thomas *et al.*, 1998; Elliott, 2001; Wang *et al.*, 2001). The bioremediation of PAH-contaminated anoxic sediments has been restricted to the application of alternative electron acceptors (Mihelcic and Luthy, 1988; Lovley *et al.*, 1989; McFarland and Sims, 1991; Lovley *et al.*, 1994; Lovley *et al.*, 1996; Coates *et al.*, 1997; Nieman *et al.*, 2001). However, little information is available on the effects of applying surfactants, inorganic or organic nutrients, or growth factors on PAH biodegradation in anoxic

* To whom correspondence should be addressed.
(Tel) 82-31-330-6691; (Fax) 82-31-336-6336
(E-mail) ysoh@mju.ac.kr

sediments.

The objective of this study was to examine the degradation of PAHs in anaerobic estuarine sediments to investigate the possible role of microbial activity. We also evaluated the effects of adding biostimulating agents (surfactants, nutrients, carbon sources, and growth factors) on the extent and rate of PAH degradation.

Materials and Methods

Sample collection

PAH-contaminated sediments were collected as grab samples from an estuary immediately adjacent to a petrochemical factory in the industrial area of Gwangyang Bay, Korea. The site was selected because of its size and the duration of hydrocarbon contamination from fuel spills and industrial wastes. We collected sediments and seawater from the site. We discarded the top 1 cm of light-brown oxic sediment and packed the remaining sulfide-blackened anoxic sediment into canning jars. The sediments and seawater were kept at 4°C during transportation




Fig. 1. Schematic representation of packed sediment in a 600 ml air-tight jar.

and were stored in the dark prior to use. The physicochemical characteristics of the sediment were determined according to standard methods (APHA, 1998) after extracting the water-soluble fractions from the sediment using deionized water. Total organic carbon was determined in terms of weight loss on ignition (LOI) at 650°C using methods described by Carter (1993).

Experimental design

We used sealed 1 L air-tight jars as packed-sediment microcosms to investigate the degradation of PAHs in the sediment (Fig. 1). Aliquots of sediment (600 ml) were packed into the jars after homogeneously mixing them in a glove box with the various treatments. Seawater collected *in situ* was added to cover the top 1-1.5 cm of sediment and then the headspace was filled with nitrogen gas. The treatments consisted of an abiotic control (AC), a no-treatment control (NTC), slow-release fertilizer (SRF), Tween 80, lactate, yeast extract (YE), SRF + Tween 80, and SRF + Tween 80 + lactate (Table 1). One gram of SRF (Chobi Ltd., Korea) consists of 148.9 mg urea-N, 18.0 mg phosphate-P, and 500 mg clay filler, with silica and latex added as support material.

At each sampling time, the overlying surface seawater was siphoned off so that the sediment could be sampled without disturbing the surface water. We sampled 10 ml sediment aliquots in triplicate using 10 ml blunt-ended syringe corers. The sediment in the jars was then topped with the same seawater again. The sediments were incubated at 20°C in the dark without shaking.

PAH extraction

The dry weight of each sediment core was measured after drying at 60°C for 10 h and finely grinding in a mortar. We added 100 µl of d₁₀-phenanthrene solution (Sigma, 200 mg d₁₀-phenanthrene in 1 L CH₂Cl₂) to the sediment as a surrogate standard. The sediments were extracted three times using 30 ml of CH₂Cl₂ in a sonic bath (Bransonic, USA) for 30 min at 30°C. The CH₂Cl₂ phase was decanted after each extraction cycle and filtered

Table 1. Treatments of PAH-contaminated sediments

Treatment	Chemical agent	Concentration (in pure water)
Abiotic control	Glutaraldehyde	5% (v/v)
Unamended control	None	
Inorganic nutrients	Slow-release fertilizer (SRF) ^a	9 mg
Surfactant	Tween 80	10x CMC ^b
Carbon source	Lactate or dextrin	1% (w/v)
Growth factor	Yeast extract	0.1% (w/v)
Surfactant + Nutrients I	Tween 80 + SRF	10x CMC ^b + 9 mg
Surfactant + Nutrients II	Tween 80 + SRF + Lactate	10x CMC ^b + 9 mg + 1% (w/v)

^a1 g SRF (Chobi Ltd., Korea) consists of 148.9 mg urea-N, 18.0 mg phosphate-P, and 500 mg clay filler, with silica and latex added as support material.

^b1x CMC, 0.075% (v/v)

through filter paper. The composite extracts were evaporated using a rotary vacuum evaporator at 30°C to <1 ml in volume. After adding 50 µl of decafluorobiphenyl (DFB) solution (200 mg DFB in 1 L CH₂Cl₂) as an internal standard, the total volume of the extract was adjusted to 1 ml with CH₂Cl₂, filtered through a PTFE 0.2 µm syringe filter (Advantec, USA), and stored in amber glass vials sealed with Teflon-butyl rubber caps. The samples were refrigerated before GC analysis.

PAH analysis

PAH concentrations in the extracts were analyzed using a Hewlett-Packard 6890 series gas chromatograph with an Agilent HP-5 fused silica capillary column (60 m × 0.32 mm i.d., film thickness 1.0 µm) and flame ionization detector (FID). Helium was used as the carrier gas at a flow rate of 1.5 ml/min and average velocity of 26 cm/sec. The temperature was ramped from 150°C to 190°C at 8°C/min with 5 min holding, then to 220°C at 2°C/min, then to 300°C at 15°C/min, and finally to 310°C at 2°C/min with 20 min holding. Each PAH was identified using PAH standard (Sigma, USA) and confirmed by GC/MS (Varian, USA) analysis.

Results and Discussion

Sediment characteristics

As shown in Table 2, the organic matter content of the Gwangyang sediments was 9.5% by dry weight. Compared to values reported for most coastal sediments (3–10%), these sediments were highly enriched in organic matter, suggesting that particulate organic inputs to Gwangyang Bay are very high (Khim *et al.*, 2001; Wang *et al.*, 2001). Total N and P concentrations were 42.5 and 0.43 µg/g dry sediment, respectively. These values are very low compared to those reported for other coastal sediments, such as those from Boston Harbor, MA, USA

Table 2. Physicochemical characteristics of Gwangyang Bay sediments

Parameters	Units	Value
pH	–	7.92
Moisture	gH ₂ O/g wet sediment	0.51
Organic matter	g/g dry sediment	0.095
Salinity	part per thousand	9.20
Total-N	µg/g dry sediment	42.50
Nitrate-N	µg/g dry sediment	31.30
Total-P	µg/g dry sediment	0.43
Phosphate-P	µg/g dry sediment	0.32
Sulfate	mg/g dry sediment	0.51
Sulfide	mg/g dry sediment	1.00
Carbonate	µmol/g dry sediment	29.10
Iron	µg/g dry sediment	18.70

(Wang *et al.*, 2001). The optimal recommended C:N:P ratio for most bioremediation applications is approximately 100:10:1 (Alexander, 1999; Maier *et al.*, 2000; Rittmann and McCarty, 2001). Our analysis showed that the sediments lacked an adequate supply of inorganic nutrients, such as nitrogen and phosphorus, for the biodegradation of PAHs. Thus supplying inorganic nutrients could be the most important step to enhance PAH biodegradation. Concentrations of sulfate, nitrate, and carbonate were not high enough to support anaerobic respiration (Table 2), which indicates that these electron acceptors can limit anaerobic respiration processes. Genthner *et al.* (1997) reported that nitrate and sulfate concentrations in the 15–20 mM range were required to biodegrade 2-ring or 3-ring PAHs in anaerobic sediments. The sediment pH of 7.92 was in the optimal range for mineralization of petroleum hydrocarbons (Gibbons, 1991; Chang *et al.*, 2002).

The concentrations of the 14 detected PAHs ranged from 9.3 to 2,889 µg/kg dry sediment (Table 3), which is significantly higher than those reported in other studies (Prah and Carpenter, 1983; Huntley and Bonnevie, 1995; Lim, 1998). The PAH distribution was characteristic of the signature of asphaltenic petroleum products, i.e., very high levels of fluoranthene, pyrene, and chrysene (Rothermich *et al.*, 2002).

Degradation of PAHs under various conditions

To evaluate the anaerobic degradation of *in situ* PAHs in

Table 3. Concentrations of PAHs in Gwangyang Bay sediment

PAH	Concentration (ng/gds ^a)
Naphthalene	N.D. ^b
Acenaphthene	60.4
Acenaphthylene	9.3
Fluorene	18.4
Phenanthrene	76.9
Anthracene	80.8
Fluoranthene	2,888.8
Pyrene	1,171.5
Benzo[<i>a</i>]anthracene	651.8
Chrysene	1,644.2
Benzo[<i>e</i>]fluoranthene	210.0
Benzo[<i>k</i>]fluoranthene	66.7
Benzo[<i>a</i>]pyrene	104.3
Indeno[1,2,3- <i>cd</i>]pyrene	N.D.
Dibenz[<i>a,h</i>]anthracene	47.8
Benzo[<i>ghi</i>]perylene	25.3
Total	7,056.2

^agds, g dry sediment

^bN.D., Not detected

the contaminated sediments, we monitored GC-detectable concentrations of PAHs for 120 days after amending with possible biostimulating agents. Compared to the PAH concentrations in the biologically active sediments, those in the sediment treated with glutaraldehyde remained virtually constant during 120 days of incubation (Table 4). This clearly demonstrates that PAH degradation is a biological process and that indigenous microorganisms in the sediment have a high potential of degrading 2- to 4-ring PAHs. Compared to the amended sediments, there was little loss of *in situ* PAHs in the unamended sediment. Additions of electron acceptors such as sulfate or carbonate also did not enhance the degradation of PAHs (data not shown). However, Tween 80, SRF, lactate, and yeast extract significantly enhanced the degradation of PAHs, which indicates that these factors were absent from or deficient in the native sediments.

Surfactants are useful for the bioremediation of environments contaminated with PAHs because they enhance the desorption and solubility of hydrophobic compounds. Many studies have sought to stimulate PAH biodegradation with surfactants that enhance the solubility and desorption rates of PAHs from soil particles under aerobic conditions (Shin *et al.*, 1999; Joo *et al.*, 2001; Garon *et al.*, 2002; Prak and Pritchard, 2002). Until recently, little attention had been paid to enhancing PAH degradation in anaerobic sediments by the use of surfactants. In this study, the addition of Tween 80 at a final concentration of 10x CMC increased the solvent extraction recovery of *in situ* PAHs from sediment samples up to 2.5 times (data not shown). The results demonstrated that when micelles were formed, PAH molecules were partitioned to their hydrophobic cores, leading to an apparent increase in the solubility of the PAHs. The increased availability of PAHs resulted in enhanced PAH biodegradation (Table 4).

A major difference between remediating aquifer and marine sediments by nutrient amendment is that dissolved nutrients can be added to groundwater, whereas nutrients should be added to marine sediments in a less soluble

form for slow release via dissolution into the sediments over several years (Adegbidi *et al.*, 2003). The addition of SRF could maintain nutrient supplements for microorganisms for a long time, as well as prevent nitrogen and phosphorus compounds from being lost by dilution at high tide (Choi *et al.*, 2003).

Although *in situ* marine sediments contain quantities of macronutrients and trace elements used by microorganisms, certain compounds essential for PAH degradation could be deficient because the microorganisms quickly use them up. We chose lactate, a readily utilizable substrate for sulfate-reducing bacteria under anaerobic conditions, and dextrin, a less favorable but long-lasting substrate, as amending agents. Yeast extract was chosen to supply essential nutrients, such as amino acids, vitamins, and trace elements, for the growth of microorganisms. As shown in Table 4, lactate and yeast extract significantly enhanced the PAH degradation rate, whereas dextrin did not. Previous studies have also reported that the presence of co-substrates such as acetate, lactate, pyruvate, yeast extract, and glucose enhance PAH degradation (Yuan *et al.*, 2000; Chang *et al.*, 2002). We predict that a preference for the type of organic nutrients exists that could be site specific.

During the 120 days of incubation, the total *in situ* PAH concentrations were determined to verify the effects of the various biostimulators on the remediation of PAH-polluted marine sediments (Fig. 2). The desorption of PAHs by the addition of Tween 80 could enrich the bioavailable PAHs for PAH-degrading microorganisms in contaminated marine sediments. The amendment of contaminated sediments with nitrogen and phosphorus in the form of SRF enhanced *in situ* PAH degradation, which indicated the significance of inorganic nutrients as limiting factors of PAH degradation in sediments. Lactate, a preferred carbon and energy source for sulfate reducers, also enhanced PAH degradation significantly. The combination of inorganic nutrients (SRF_s), a surfactant (Tween 80), and an organic nutrient (lactate) increased PAH degradation rates more than did the individual compounds

Table 4. Effects of biostimulating agents* on PAH degradation in sediments

PAH	Degradation rate of PAHs (µg-PAH/kg-dry sediment/day)						
	GAL	NTC	T80	SRF	LAC	DEX	YE
Phenanthrene	0.0	0.0	1.3	2.0	2.1	0.0	1.8
Anthrathene	0.0	0.3	2.9	3.5	5.1	0.0	0.2
Fluoranthene	0.0	0.8	1.6	0.7	11.1	0.2	3.8
Pyrene	0.0	0.0	0.4	0.1	7.5	0.5	1.6
Chrysene	0.0	0.8	4.4	2.1	2.4	0.7	2.3
Benzo[a]pyrene	0.0	1.7	2.1	6.1	1.4	0.0	0.0
Total	0.0	3.6	12.7	14.5	29.6	1.4	9.7

*GAL, glutaraldehyde abiotic control; NTC, unamended control; T80, Tween 80; SRF, slow-release fertilizer; LAC, lactate; DEX, dextrin; YE, yeast extract. Table 1 lists the concentration of each agent.




Fig. 2. The time course of PAH degradation in sediment with various treatments. ●, control; ○, SRF; ▲, Tween 80; ■, lactate; □, SRF + Tween 80 + lactate.

(Fig. 2). Thus, amending the PAH-contaminated sediment with biostimulators increased the intrinsic biodegradation potentials of these PAHs. Our results will contribute to the development of new strategies for *in situ* treatment of PAH-contaminated anoxic sediments in Gwangyang Bay.

Acknowledgment

This work was supported by the Ecotechnopia program of the Ministry of Environment, Republic of Korea.

References

- Adegbidi, H.G., R.D. Briggs, T.A. Volk, E.H. White, and L.P. Abrahamson. 2003. Effect of organic amendments and slow-release nitrogen fertilizer on willow biomass production and soil chemical characteristics. *Biomass and Bioenergy* 25, 389-398.
- Alexander, M. 1999. Biodegradation and bioremediation, 2nd ed. p. 275. Academic Press, San Diego, California.
- APHA, AWWA, WEF. 1998. Standard methods for the examination of water and wastewater, 20th ed. APHA, Washington, DC.
- Carter, M.R. 1993. Soil sampling and methods of analysis, p. 141-394. Canadian Society of Soil Science, Lewis Publishers, Boca Raton, Florida.
- Chang, B.V., S.W. Chang, and S.Y. Yuan. 2003. Anaerobic degradation of polycyclic aromatic hydrocarbons in sludge. *Adv. Environ. Res.* 7, 623-628.
- Chang, B.V., L.C. Shiung, and S.Y. Yuan. 2002. Anaerobic biodegradation of polycyclic aromatic hydrocarbon in soil. *Chemosphere* 48, 717-724.
- Choi, S.-C., Y.-H. Lee, and Y.-S. Oh. 2003. Treatability tests for the bioremediation of unsanitary landfill waste soil. *J. Microbiol.* 41, 169-173.
- Coates, J.D., J. Woodward, J. Allen, P. Philp, and D.R. Lovley. 1997. Anaerobic degradation of polycyclic aromatic hydrocarbons and alkanes in petroleum-contaminated marine harbor sediments. *Appl. Environ. Microbiol.* 63, 3589-3593.

- Elliott, M. 2001. Polycyclic aromatic hydrocarbons and redox parameter in a creosote-contaminated aquifer. MS thesis. Virginia Polytechnic Institute and State University, USA.
- Garon, D., S. Krivobok, D. Wouessidjewe, and F. Seigle-Murandi. 2002. Influence of surfactant on solubilization and fungal degradation of fluorene. *Chemosphere* 47, 303-309.
- Gibbons, J.H. 1991. Bioremediation for marine oil spills. U.S. Congress, Office of Technology Assessment. Washington D.C.
- Genthner, B.R.S., G.T. Townsend, S.E. Lantz, and J.G. Mueller. 1997. Persistence of polycyclic aromatic hydrocarbon components of creosote under anaerobic enrichment conditions. *Arch. Environ. Contam. Toxicol.* 32, 99-105.
- Han, M.-J., H.-T. Choi, and H.-G. Song. 2003. Degradation of phenanthrene by *Trametes versicolor* and its laccase. *J. Microbiol.* 42, 94-98.
- Hayes, L.A., K.P. Nevin, and D.R. Lovley. 1999. Role of prior exposure on anaerobic degradation of naphthalene and phenanthrene in marine harbor sediments. *Organic Geochem.* 30, 937-945.
- Huntley, S.L. and N.L. Bonnevie. 1995. Polycyclic aromatic hydrocarbon and petroleum hydrocarbon contamination in sediment from the Newark Bay Estuary, New Jersey. *Arch. Environ. Contam. Toxicol.* 28, 93-107.
- Joo, C.-S., Y.-S. Oh, and W.-J. Chung. 2001. Evaluation of bioremediation effectiveness by resolving rate-limiting parameters in diesel-contaminated soil. *J. Microbiol. Biotechnol.* 11, 607-613.
- Kanally, R.A. and S. Harayama. 2000. Biodegradation of high-molecular-weight polycyclic aromatic hydrocarbons by bacteria. *J. Bacteriol.* 182, 2059-2067.
- Karthikeyan, R. and A. Bhandari. 2001. Anaerobic biotransformation of aromatic and polycyclic aromatic hydrocarbons in soil microcosms: a review. *Hazard. Subst. Res.* 3, 1-19.
- Khim, J.S., K.T. Lee, K. Kannan, D.L. Villeneuve, J.P. Giesy, and C.H. Koh. 2001. Trace organic contaminants in sediment and water from Ulsan Bay and this vicinity, Korea. *Arch. Environ. Contam. Toxicol.* 40, 141-150.
- Lim, W.-H. 1998. Contamination of polycyclic aromatic hydrocarbons (PAHs) in Masan Bay, Korea. MS thesis. Seoul National University, Korea.
- Lovley, D.R., M.J. Baedeker, D.J. Lonergan, I.M. Cozzarelli, E.J.P. Phillips, and D.I. Siegel. 1989. Oxidation of aromatic contaminants coupled to microbial iron reduction. *Nature* 339, 297-300.
- Lovley, D.R., J.C. Woodward, and F.H. Chapelle. 1994. Stimulated anoxic biodegradation of aromatic hydrocarbons using Fe(III) ligands. *Nature* 370, 128-131.
- Lovley, D.R., J.C. Woodward, and F.H. Chapelle. 1996. Rapid anaerobic benzene oxidation with a variety of chelated Fe(III) forms. *Appl. Environ. Microbiol.* 62, 288-291.
- Maier, R.M., I.L. Pepper, and C.P. Gerba. 2000. Environmental microbiology, p. 335. Academic Press. San Diego, California.
- McFarland, M.J. and R.C. Sims. 1991. Thermodynamic framework for evaluating PAH degradation in the subsurface. *Groundwater* 29, 885-896.
- Mihelcic, J.R. and R.G. Luthy. 1988. Degradation of polycyclic aromatic hydrocarbon compounds under various redox conditions in soil-water systems. *Appl. Environ. Microbiol.* 54, 1182-1187.
- Nieman, J.K.C., R.C. Sims, J.E. McLean, J.L. Sims, and D.L. Sorensen. 2001. Fate of pyrene in contaminated soil amended

- with alternate electron acceptors. *Chemosphere* 44, 1265-1271.
- Prahl, F.G. and R. Carpenter. 1983. Polycyclic aromatic hydrocarbon-phase associations in Washington coastal sediment. *Geochim. Cosmochim. Acta* 47, 1013-1023.
- Prak, D.J.L. and P.H. Pritchard. 2002. Solubilization of polycyclic aromatic hydrocarbon mixtures in micellar nonionic surfactant solutions. *Wat. Res.* 36, 3463-3472.
- Rittmann, B.E. and P.L. McCarty. 2001. Environmental biotechnology: principles and applications, p. 535-537. McGraw-Hill, New York, New York.
- Rothermich, M.M, L.A. Hayes, and D.R. Lovley. 2002. Anaerobic, sulfate-dependent degradation of polycyclic aromatic hydrocarbons in petroleum-contaminated harbor sediment. *Environ. Sci. Technol.* 36, 4811-4817.
- Shin, S.-K., Y.-S. Oh, and S.-J. Kim. 1999. Biodegradation of phenanthrene by *Sphingomonas* sp. strain KH3-2. *J. Microbiol.* 37, 185-192.
- Thomas, S.P., H.D. Stensel, and S.E. Strand. 1998. Biodegradation of polyaromatic hydrocarbons by marine bacteria: effect of solid phase on degradation kinetics. *Wat. Res.* 33, 868-880.
- Wang, X.-C., Y.-X. Zhang, and R.F. Chen. 2001. Distribution and partitioning of polycyclic aromatic hydrocarbons (PAHs) in different size fractions in sediments from Boston Harbor, United States. *Mar. Pollut. Bull.* 42, 1139-1149.
- Weiner, J.M. and D.R. Lovley. 1998. Rapid benzene degradation in methanogenic sediments from a petroleum-contaminated aquifer. *Appl. Environ. Microbiol.* 64, 1937-1939.
- Yuan, S.Y., S.H. Wei, and B.V. Chang. 2000. Biodegradation of polycyclic aromatic hydrocarbons by a mixed culture. *Chemosphere* 41, 1463-1468.